



ELSEVIER

Discrete Mathematics 263 (2003) 295–296

DISCRETE
MATHEMATICS

www.elsevier.com/locate/disc

Note

Incremental construction properties in dimension two—shellability, extendable shellability and vertex decomposability[☆]

Sonoko Moriyama, Fumihiko Takeuchi*

Department of Information Science, University of Tokyo, Tokyo 113-0033, Japan

Received 30 November 2000; received in revised form 4 September 2001; accepted 24 September 2001

Abstract

We give new examples of shellable, but not extendably shellable two-dimensional simplicial complexes. They include minimal examples that are smaller than those previously known. We also give new examples of shellable, but not vertex decomposable two-dimensional simplicial complexes, including extendably shellable ones. This shows that neither extendable shellability nor vertex decomposability implies the other. We found these examples by enumerating shellable two-dimensional simplicial complexes that are not pseudomanifolds.

© 2002 Elsevier Science B.V. All rights reserved.

Keywords: Incremental construction; Shellability; Extendable shellability; Vertex decomposability; Combinatorial topology; Enumeration

A pure n -dimensional simplicial complex \mathcal{X} —or, for short, an n -complex—is *shellable* if its facets (i.e., n -simplices) can be ordered linearly so that the intersection of any facet with the union of its preceding facets (relative to that order) is a union of $(n - 1)$ -simplices, and any such linear order is called a *shelling* of \mathcal{X} (see [1,4] or [5] for details and examples). An n -complex \mathcal{X} is *extendably shellable* if any shelling of an n -subcomplex of \mathcal{X} can be continued to a shelling of \mathcal{X} .

Consider the following five 2-complexes defined in terms of a complete list of their facets (see [5] for further examples). In the table, 1-simplices that are contained in one facet, only, are indicated by bold letters. For shellable, but not extendably shellable

[☆] Preliminary version appeared in Proceedings of the 12th Canadian Conference on Computational Geometry, Fredericton, 2000, pp. 65–72. Electronic Annexes available. See <http://www.elsevier.com/locate/disc>.

* Corresponding author.

E-mail addresses: moriso@is.s.u-tokyo.ac.jp (S. Moriyama), fumi@is.s.u-tokyo.ac.jp (F. Takeuchi).

complexes, the stuck partial shellings (unsorted, as sets, which happen to be unique in these examples) are also provided:

V6F9-1	124, 126 , 134, 135 , 245, 256, 346, 356, 456 stuck partial shelling: {124, 126 , 134, 135 }
V6F9-2	123, 126 , 135 , 234, 245, 256, 346, 356, 456 stuck partial shelling: {123, 126 , 135 , 234}
V6F10-1[3]	123 , 124, 126 , 134, 135 , 245, 256, 346, 356, 456 stuck partial shelling: { 123 , 124, 126 , 134, 135 }
V6F10-6	123 , 124, 125 , 134, 136 , 245, 256 , 346, 356 , 456
V6F10-7	123, 125, 126 , 134, 145, 234 , 256, 346, 356 , 456

Theorem A. *The two complexes V6F9-1,2 having 6 vertices and 9 facets are shellable, but not extendably shellable. There is no 2-complex with less than 6 vertices or less than 9 facets having this property.*

An n -complex \mathcal{X} is *vertex decomposable* if it has only one facet, or if it has a vertex v such that both, its *link* $\text{link}_{\mathcal{X}}(v) := \{X - \{v\} \mid v \in X \in \mathcal{X}\}$ as well as its *deletion* $\text{del}_{\mathcal{X}}(v) := \{X \in \mathcal{X} \mid v \notin X\}$, are vertex decomposable. Vertex decomposability is known to imply shellability.

Theorem B. *The three complexes V6F10-1,6,7 having 6 vertices and 10 facets are shellable, but not vertex decomposable. There is no 2-complex with less than 6 vertices or with 6 vertices and less than 10 facets having this property. Furthermore, V6F10-1 is not extendably shellable while V6F10-6,7 are extendably shellable.*

Vertex decomposable, yet not extendably shellable complexes have been known while our examples of extendably shellable, yet not vertex decomposable complexes appear to be new. Together, these examples demonstrate that none of these two stronger properties implies other.

Acknowledgements

This paper is based on [2]. The authors thank David Avis, Hiroshi Imai, and especially Masahiro Hachimori for discussions and encouragement.

References

- [1] A. Björner, Topological methods, in: R.L. Graham, M. Grötschel, L. Lovász, (Eds.), Handbook of Combinatorics, Elsevier, Amsterdam, 1995, pp. 1819–1872.
- [2] S. Moriyama, Extendable shellability in dimension two, Bachelor Thesis, University of Tokyo, Tokyo, 2000.
- [3] R.S. Simon, Combinatorial properties of cleanness, J. Algebra 167 (1994) 361–388.
- [4] G.M. Ziegler, Lectures on Polytopes, Springer, New York, 1995.
- [5] Electronic Annexes available. See <http://www.elsevier.com/locate/disc>.